

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 31/495, 9/20</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/13687</b>
			(43) International Publication Date: 16 March 2000 (16.03.00)
(21) International Application Number: <b>PCT/US99/20968</b>			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 9 September 1999 (09.09.99)			
(30) Priority Data: 60/099,804 10 September 1998 (10.09.98) US 09/321,522 27 May 1999 (27.05.99) US			
(71) Applicant (for all designated States except US): CV THERAPEUTICS, INC. [US/US]; 3172 Porter Drive, Palo Alto, CA 94304 (US).			
(72) Inventor; and (75) Inventor/Applicant (for US only): WOLFF, Andrew, A. [US/US]; 1140 Lawton Street, San Francisco, CA 94122 (US).			
(74) Agent: HUGHES, A., Blair; McDonnell, Boehnen, Hulbert & Berghoff, Suite 3200, 300 S. Wacker Drive, Chicago, IL 60606 (US).			Published Without international search report and to be republished upon receipt of that report.
(54) Title: SUSTAINED RELEASE RANOLAZINE FORMULATIONS			
(57) Abstract <p>A sustained release ranolazine formulation contains an intimate mixture of ranolazine and a partially neutralized pH-dependent binder to form a film that is mostly insoluble in aqueous media below pH 4.5 and soluble in aqueous media above pH 4.5. The formulation is suitable for twice daily administration of ranolazine and is useful for controlling the rate of dissolution of ranolazine, and to maintain human plasma ranolazine levels at between 550 and 7500 ng base/mL.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MR	Mauritania	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MW	Malawi	UA	Ukraine
BR	Brazil	IL	Israel	MX	Mexico	UG	Uganda
BY	Belarus	IS	Iceland	NE	Niger	US	United States of America
CA	Canada	IT	Italy	NL	Netherlands	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NO	Norway	VN	Viet Nam
CG	Congo	KE	Kenya	NZ	New Zealand	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	PL	Poland	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PT	Portugal		
CM	Cameroon	KR	Republic of Korea	RO	Romania		
CN	China	KZ	Kazakhstan	RU	Russian Federation		
CU	Cuba	LC	Saint Lucia	SD	Sudan		
CZ	Czech Republic	LI	Liechtenstein	SE	Sweden		
DE	Germany	LK	Sri Lanka	SG	Singapore		
DK	Denmark	LR	Liberia				
EE	Estonia						

## SPECIFICATION

**TITLE: SUSTAINED RELEASE RANOLAZINE FORMULATIONS**

5

**BACKGROUND OF THE INVENTION**

This application claims priority to U.S. Provisional Patent Application serial number 60/099804 filed on September 10, 1998.

10    **1. Field of the Invention**

This invention relates to a method for using an orally administered sustained release ranolazine dosage formulations to maintain human ranolazine plasma levels at therapeutic levels.

**2. Description of the Art**

15       U.S. Patent No. 4,567,264, the specification of which is incorporated herein by reference, discloses ranolazine, (±)-N- (2,6-dimethylphenyl)-4-[2-hydroxy-3- (2-methoxyphenoxy)-propyl]-1-piperazineacetamide, and its pharmaceutically acceptable salts, and their use in the treatment of cardiovascular diseases, including arrhythmias, variant and exercise-induced angina, and myocardial infarction.

20       U.S. Patent No. 5,506,229, which is incorporated herein by reference, discloses the use of ranolazine and its pharmaceutically acceptable salts and esters for the treatment of tissues experiencing a physical or chemical insult, including cardioplegia, hypoxic or reperfusion injury to cardiac or skeletal muscle or brain tissue, and for use in transplants. Conventional oral and parenteral formulations are disclosed, including controlled release formulations. In  
25   particular, Example 7D of U.S. Patent No. 5,506,229 describes a controlled release formulation in capsule form comprising microspheres of ranolazine and microcrystalline cellulose coated with release controlling polymers.

The presently preferred route of administration for ranolazine and its pharmaceutically acceptable salts and esters is oral. A typical oral dosage form is a compressed tablet, a hard gelatin capsule filled with a powder mix or granulate, or a soft gelatin capsule (softgel) filled with a solution or suspension. U.S. Patent No. 5,472,707, the specification of which is  
5 incorporated herein by reference, discloses a high-dose oral formulation employing supercooled liquid ranolazine as a fill solution for a hard gelatin capsule or softgel.

As set forth in application Example 3, the initial trials of ranolazine on humans suffering from angina were failures. The trials used an immediate release of ranolazine formulation at a dose level of 120mg taken three times daily. Based upon the initial  
10 experiments, it was uncertain whether or not ranolazine could be given to humans in an amount and mode that is effective against angina.

One problem with conventional oral dosage formulations is that they are not ideally suited to ranolazine and its pharmaceutically acceptable salts, because the solubility of ranolazine is relatively high at the low pH that occurs in the stomach. Furthermore ranolazine  
15 also has a relatively short plasma half-life. The high acid solubility property of ranolazine results in rapid drug absorption and clearance, causing large and undesirable fluctuations in plasma concentration of ranolazine and a short duration of action, thus necessitating frequent oral administration for adequate treatment.

There is therefore a need for a method for administering ranolazine in an oral dosage  
20 form once or twice daily that provides therapeutically effective plasma concentrations of ranolazine for the treatment of angina in humans.

**SUMMARY OF THE INVENTION**

In a first aspect, this invention is a sustained release ranolazine formulation where the majority of the formulation consists of active ranolazine.

In another aspect, this invention is a method of treating a human patient who is  
5 suffering from angina or other coronary disorders by administering a sustained release ranolazine formulation to the patient once or twice daily.

In yet another aspect, this invention is a method for treating a mammal having a disease state for which the administration of ranolazine is indicated, comprising the once or twice-daily administration of sustained release ranolazine formulation of this invention in a  
10 manner that maintains plasma ranolazine close to minimal effective levels without peak fluctuations.

Still another aspect of this invention is a method of maintaining useful levels of ranolazine in human plasma by administering a ranolazine containing dosage form only one or twice daily.

15 This invention includes methods for treating a human patient suffering from a cardiovascular disease selected from arrhythmias, variant and exercise-induced angina, and myocardial infarction. The method includes administering a sustained release pharmaceutical dosage form including at least 50% by weight ranolazine in no more than two tablets per dose to the human patient to maintain ranolazine plasma levels in the human patient of from about  
20 550 to about 7500 ng base/mL for at least 24 hours wherein the dose is administered at a frequency selected from once, twice and three times over 24 hours.

This invention further includes methods for treating a human patient suffering from a cardiovascular disease selected from arrhythmias, variant and exercise-induced angina, and myocardial infarction. The method includes administering a sustained release pharmaceutical  
25 dosage form including from about 70 to about 80% by weight ranolazine in no more than two

tablets per dose to the human patient to maintain ranolazine plasma levels in the human patient of from about 1000 to about 3900 ng base/mL for at least 24 hours wherein the dose is administered at a frequency selected from one and two times over 24 hours.

This invention also includes pharmaceutical dosage forms comprising at least about 50  
5 wt% ranolazine and at least one pH dependent binder that inhibits the release of ranolazine from the sustained release dosage form when the sustained release dosage form is subjected to an aqueous environment having a pH of the stomach and that promotes the release of a therapeutic amount of ranolazine in an aqueous solution having a pH above about 4.5.

10

## DETAILED DESCRIPTION OF THE INVENTION

"Ranolazine" is the compound ( $\pm$ )-N- (2,6-dimethylphenyl)-4-[2-hydroxy-3- (2-methoxyphenoxy) propyl]-1-piperazine-acetamide, or its enantiomers (R)-(+)-N-(2,6-dimethylphenyl)-4-[2-hydroxy-3-(2-methoxyphenoxy)-propyl]-1-piperazineacetamide, and  
5 (S)-(-)-N-(2,6-dimethylphenyl) -4-[2-hydroxy-3-(2-methoxyphenoxy)-propyl]-1-piperazineacetamide and their pharmaceutically acceptable salts, and mixtures thereof. Unless otherwise stated the ranolazine plasma concentrations used in the specification and examples refers to ranolazine free base.

"Optional" and "optionally" mean that the subsequently described event or  
10 circumstance may or may not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optional pharmaceutical excipients" indicates that a formulation so described may or may not include pharmaceutical excipients other than those specifically stated to be present, and that the formulation so described includes instances in which the optional excipients are present and  
15 instances in which they are not.

"Treating" and "treatment" refer to any treatment of a disease in a mammal, particularly a human, and include:

- (i) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it;
- 20 (ii) inhibiting the disease, i.e., arresting its development; or
- (iii) relieving the disease, i.e., causing regression of the disease.

"Immediate release" ("IR") refers to formulations or dosage units that rapidly dissolve *in vitro* and are intended to be completely dissolved and absorbed in the stomach or upper gastrointestinal tract. Conventionally, such formulations release at least 90% of the active  
25 ingredient within 30 minutes of administration.

"Sustained release" ("SR") refers to formulations or dosage units of this invention that are slowly and continuously dissolved and absorbed in the stomach and gastrointestinal tract over a period of about six hours or more. Preferred sustained release formulations are those exhibiting plasma concentrations of ranolazine suitable for no more than twice daily administration with two or less tablets per dosing as described below.

Plasma ranolazine concentration is a mean concentration determined by analyzing the concentration of ranolazine in as few as five to as many as ten humans who are on the same dosing schedule. It is important that the ranolazine concentration is a mean value because of variances in ranolazine concentrations in individuals that may be caused by differences in weight, metabolism, or disease states which may cause one person to metabolize ranolazine faster or slower than an average person. The plasma ranolazine levels are determined from drawn blood onto heparin.

Definitions of other terms used in this application are:

ANOVA = analysis of variance

ATP = adenosine triphosphate

ECG = electrocardiographic

ETT = exercise treadmill test

PDH = pyruvate dehydrogenase

$C_{max}$  = maximum concentration

$C_{trough}$  = residual concentration at 8 hours post-dose for IR formulations and 12 hours post-dose for SR formulations A-C of Example 2.

tid = three times per day

bid = twice daily

$C_x$  = concentration at time x

$T_{max}$  = time to maximum concentration

$AUC_x$  = area under the curve after x hours or time interval



Percentages given are percentages by weight, unless otherwise stated. This invention involves sustained release ranolazine dosage forms as well as methods for administering sustained release ranolazine dosage forms of this invention to provide for therapeutic plasma levels of ranolazine.

5       The sustained release ranolazine formulations of this invention are preferably in the form of a compressed tablet comprising an intimate mixture of ranolazine and a partially neutralized pH-dependent binder that controls the rate of ranolazine dissolution in aqueous media across the range of pH in the stomach (typically approximately 2) and in the intestine (typically approximately about 5.5).

10       To provide for a sustained release of ranolazine, one or more pH-dependent binders are chosen to control the dissolution profile of the ranolazine formulation so that the formulation releases ranolazine slowly and continuously as the formulation passed through the stomach and gastrointestinal tract. The dissolution control capacity of the pH-dependent binder(s) is particularly important in a sustained release ranolazine formulation because a  
15       sustained release formulation that contains sufficient ranolazine for twice daily administration may cause untoward side effects if the ranolazine is released too rapidly ("dose-dumping").

      Accordingly, the pH-dependent binders suitable for use in this invention are those which inhibit rapid release of drug from a tablet during its residence in the stomach (where the pH is below about 4.5), and which promotes the release of a therapeutic amount of ranolazine  
20       from the dosage form in the lower gastrointestinal tract (where the pH is generally greater than about 4.5). Many materials known in the pharmaceutical art as "enteric" binders and coating agents have the desired pH dissolution properties. These include phthalic acid derivatives such as the phthalic acid derivatives of vinyl polymers and copolymers, hydroxyalkylcelluloses, alkylcelluloses, cellulose acetates, hydroxyalkylcellulose acetates,  
25       cellulose ethers, alkylcellulose acetates, and the partial esters thereof, and polymers and

copolymers of lower alkyl acrylic acids and lower alkyl acrylates, and the partial esters thereof.

Preferred pH-dependent binder materials which can be used in conjunction with ranolazine to create a sustained release formulation are methacrylic acid copolymers.

5 Methacrylic acid copolymers are copolymers of methacrylic acid with neutral acrylate or methacrylate esters such as ethyl acrylate or methyl methacrylate. A most preferred copolymer is methacrylic acid copolymer, Type C, USP (which is a copolymer of methacrylic acid and ethyl acrylate having between 46.0% and 50.6% methacrylic acid units). Such a copolymer is commercially available, from Röhm Pharma as Eudragit® L 100-55 (as a

10 powder) or L30D-55 (as a 30% dispersion in water). Other pH-dependent binder materials which may be used alone or in combination in a sustained release ranolazine dosage form include hydroxypropyl cellulose phthalate, hydroxypropyl methylcellulose phthalate, cellulose acetate phthalate, polyvinylacetate phthalate, polyvinylpyrrolidone phthalate, and the like. One or more pH-dependent binders are present in the ranolazine dosage forms of this

15 invention in an amount ranging from about 1 to about 20 wt%, more preferably from about 5 to about 12 wt% and most preferably about 10 wt%.

One or more pH-independent binders may be in used in sustained release ranolazine oral dosage forms. It is to be noted that pH-dependent binders and viscosity enhancing agents such as hydroxypropyl methylcellulose, hydroxypropyl cellulose, methylcellulose,

20 polyvinylpyrrolidone, neutral poly(meth)acrylate esters, and the like, do not themselves provide the required dissolution control provided by the identified pH-dependent binders. The pH-independent binders are present in the formulation of this invention in an amount ranging from about 1 to about 10 wt%, and preferably in amount ranging from about 1 to about 3 wt% and most preferably about 2.0 wt%.

As shown in Table 1, ranolazine is relatively insoluble in aqueous solutions having a pH above about 6.5, while the solubility begins to increase dramatically below about pH 6.

Table 1

Solution pH	Solubility (mg/mL)	USP Solubility Class
4.81	161	Freely soluble
4.89	73.8	Soluble
4.90	76.4	Soluble
5.04	49.4	Soluble
5.35	16.7	Sparingly soluble
5.82	5.48	Slightly soluble
6.46	1.63	Slightly soluble
6.73	0.83	Very slightly soluble
7.08	0.39	Very slightly soluble
7.59 (unbuffered water)	0.24	Very slightly soluble
7.73	0.17	Very slightly soluble
12.66	0.18	Very slightly soluble

5 Increasing the pH-dependent binder content in the formulation decreases the release rate of ranolazine from the formulation at pH is below 4.5 typical of the pH found in the stomach. The enteric coating formed by the binder is less soluble and increases the relative release rate above pH 4.5, where the solubility of ranolazine is lower. A proper selection of the pH-dependent binder allows for a quicker release rate of ranolazine from the formulation above

10 pH 4.5, while greatly affecting the release rate at low pH. Partial neutralization of the binder facilitates the conversion of the binder into a latex like film which forms around the individual ranolazine granules. Accordingly, the type and the quantity of the pH-dependent binder and amount of the partial neutralization composition are chosen to closely control the rate of dissolution of the ranolazine from the formulation.

15 The dosage forms of this invention should have a quantity of pH-dependent binders sufficient to produce a sustained release formulation from which the release rate of ranolazine is controlled such that at low pHs (below about 4.5) the rate of dissolution is significantly slowed. In the case of methacrylic acid copolymer, type C, USP (Eudragit® L 100-55), a suitable quantity of pH-dependent binder is between 5% and 15%. The pH dependent binder

will typically have from about 1 to about 20% of the binder methacrylic acid carboxyl groups neutralized. However, it is preferred that the degree of neutralization ranges from about 3 to 6%.

The sustained release formulation may also contain pharmaceutical excipients intimately admixed with the ranolazine and the pH-dependent binder. Pharmaceutically acceptable excipients may include, for example, pH-independent binders or film-forming agents such as hydroxypropyl methylcellulose, hydroxypropyl cellulose, methylcellulose, polyvinylpyrrolidone, neutral poly(meth)acrylate esters (e.g. the methyl methacrylate/ethyl acrylate copolymers sold under the trademark Eudragit® NE by Röhm Pharnal, starch, gelatin, sugars, carboxymethylcellulose, and the like. Other useful pharmaceutical excipients include diluents such as lactose, mannitol, dry starch, microcrystalline cellulose and the like; surface active agents such as polyoxyethylene sorbitan esters, sorbitan esters and the like; and coloring agents and flavoring agents. Lubricants (such as talc and magnesium stearate) and other tableting aids are also optionally present.

The sustained release ranolazine formulations of this invention have a ranolazine content of above about 50% by weight to about 95% or more by weight, more preferably between about 70% to about 90% by weight and most preferably from about 70 to about 80% by weight; a pH-dependent binder content of between 5% and 40%, preferably between 5% and 25%, and more preferably between 5% and 15%; with the remainder of the dosage form comprising pH-independent binders, fillers, and other optional excipients.

Particularly preferred sustained release ranolazine formulations of this invention consist, essentially of:

<u>Ingredient</u>	<u>Weight Range (%)</u>	<u>Preferred Range (%)</u>	<u>Most Preferred (%)</u>
-------------------	-----------------------------	--------------------------------	---------------------------

	Ranolazine	50-95	70-90	75
	Microcrystalline cellulose (filler)	1-35	5-15	10.6
	Methacrylic acid copolymer	1-35	5-12.5	10.0
5	Sodium hydroxide	0.1-1.0	0.2-0.6	0.4
	Hydroxypropyl methylcellulose	0.5-5.0	1-3	2.0
	Magnesium stearate	0.5-5.0	1-3	2.0

The sustained release ranolazine formulations of this invention are prepared as follows: ranolazine and pH-dependent binder and any optional excipients are intimately mixed (dry-blended). The dry-blended mixture is then granulated in the presence of an aqueous solution of a strong base which is sprayed into the blended powder. The granulate is dried, screened, mixed with optional lubricants (such as talc or magnesium stearate), and compressed into tablets. Preferred aqueous solutions of strong bases are solutions of alkali metal hydroxides, such as sodium or potassium hydroxide, preferably sodium hydroxide, in water (optionally containing up to 25% of water-miscible solvents such as lower alcohols).

The resulting ranolazine containing tablets may be coated with an optional film-forming agent, for identification, taste-masking purposes and to improve ease of swallowing. The film forming agent will typically be present in an amount ranging from between 2% and 4% of the tablet weight. Suitable film-forming agents are well-known to the art and include hydroxypropyl methylcellulose, cationic methacrylate copolymers (dimethylaminoethyl methacrylate/ methyl-butyl methacrylate copolymers - Eudragit® E - Röhm Pharma), and the like. These film-forming agents may optionally contain colorants, plasticizers, and other supplemental ingredients.

The compressed tablets preferably have a hardness sufficient to withstand 8 Kp compression. The tablet size will depend primarily upon the amount of ranolazine in the tablet. The tablets will include from 300 to 1100 mg of ranolazine free base. Preferably, the tablets will include amounts of ranolazine free base ranging from 400-600 mg, 650-850 mg,  
5 and 900-1100 mg.

In order to influence the dissolution rate, the time during which the ranolazine containing powder is wet mixed is controlled. Preferably the total powder mix time, *i.e.* the time during which the powder is exposed to sodium hydroxide solution, will range from 1 to 10 minutes and preferably from 2 to 5 minutes. Following granulation, the particles are  
10 removed from the granulator and placed in a fluid bed dryer for drying at about 60°C.

Surprisingly, it has been found that these methods produce sustained release ranolazine formulations that provide lower peak plasma ranolazine levels and yet effective plasma concentrations of ranolazine for up to 12 hours and more after administration, when the ranolazine used as its free base, rather than as the more pharmaceutically common  
15 ranolazine dihydrochloride salt or as another salt or ester. The use of ranolazine free base affords at least one advantage: The proportion of ranolazine in the tablet can be increased, since the molecular weight of ranolazine free base is only 85% that of ranolazine dihydrochloride. In this manner, delivery of an effective amount of ranolazine is achieved while limiting the physical size of the dosage unit.

20 Another advantage of sustained release ranolazine formulations of this invention is that they are prepared by a process that essentially involves only water as a solvent, and utilizes standard pharmaceutical processing techniques and equipment.

The sustained release ranolazine formulations of this invention can be used for treating cardiovascular diseases, including arrhythmias, variant and exercise-induced angina, and  
25 myocardial infarction; treatment of tissues experiencing a physical or chemical insult,

including cardioplegia, hypoxic or reperfusion injury to cardiac or skeletal muscle or brain tissue, and ischemia; and peripheral arterial diseases, such as intermittent claudication. It is most preferred that the sustained release dosage formulation be used as a mammalian anti-anginal agent and most preferably as a human anti-anginal agent.

- 5           The oral sustained release ranolazine dosage formulations of this invention are administered one, twice; or three times in a 24 hour period in order to maintain a plasma ranolazine level above the threshold therapeutic level and below the maximally tolerated levels, of between about 550 and 7500 ng base/mL in a patient. This corresponds to an amount of ranolazine 2 HCl ranging from about 644 ng/mL to about 8782 ng/mL.
- 10   Furthermore, the timing of the oral ingestion of the ranolazine oral dosage forms should be controlled to insure that the plasma ranolazine level does not exceed about 7500 ng base/mL and preferably so that the plasma ranolazine level does not exceed about 5000 ng base/mL and most preferably so that it does not exceed 3800 ng base/mL. In some instances it may be beneficial to limit the peak plasma ranolazine level to no more than about ng base/mL. At the
- 15   same time, the plasma trough ranolazine levels should preferably not fall below about 1000 ng base/mL, and in some instances should not fall below 1700 ng base/mL.

In order to achieve the preferred plasma ranolazine level of from about 1000 to about 3800 ng base/mL, it is preferred that the oral ranolazine dosage forms described herein are administered once or twice daily. If the dosage forms are administered twice daily, then it is

20   preferred that the oral ranolazine dosage forms are administered at about twelve hour intervals.

In addition to formulating and administering oral sustained release dosage forms of this invention in a manner that controls the plasma ranolazine levels, it is also important to minimize the difference between peak and trough plasma ranolazine levels. The peak plasma

25   ranolazine levels are typically achieved at from about 30 minutes to eight hours or more after

initially ingesting the dosage form while trough plasma ranolazine levels are achieved at about the time of ingestion of the next scheduled dosage form. It is preferred that the sustained release dosage forms of this invention are administered in a manner that allows for a peak ranolazine level no more than 8 times greater than the trough ranolazine level, preferably no  
5 more than 4 times greater than the trough ranolazine and most preferably no greater than 2 times trough ranolazine level.

The sustained release ranolazine formulations of this invention provide the therapeutic advantage of minimizing variations in ranolazine plasma concentration while permitting, at most, twice-daily administration. The formulation may be administered alone, or (at least  
10 initially) in combination with an immediate release formulation if rapid achievement of a therapeutically effective plasma concentration of ranolazine is desired or by soluble IV formulations and oral dosage forms.

The following Examples are representative of the invention, but are not to be construed as limiting the scope of the claims.



**EXAMPLES**

These Examples detail methods for manufacturing ranolazine dosage forms as well as experiments performed to evaluate the effectiveness of ranolazine administration and effectiveness. Throughout these Examples it should be noted that:

5       (1) Oral doses of the instant release (IR) formulation were given as capsules or tablets of the dihydrochloride salt and are expressed as the dihydrochloride salt.

(2) Oral doses of the sustained release (SR) formulation were given as tablets of the ranolazine base and are expressed as the base.

(3) When IR and SR formulations were compared in the same study, doses are  
10 expressed in terms of both base and dihydrochloride. The conversion factor for dihydrochloride to base is 0.854 (e.g.: 400 mg dihydrochloride x 0.854 = 342 mg free base equivalent).

(4) All plasma levels and pharmacokinetic parameters are expressed as levels of free base.

**EXAMPLE 1**

This Example describes a method of preparing immediate release (IR) ranolazine formulations. Ranolazine dihydrochloride (4000 g), microcrystalline cellulose (650 g), polyvinylpyrrolidone (100 g), and croscarmellose sodium (100 g) powders were intimately  
5 mixed together in a Fielder PMA 65 mixer-granulator, and sufficient water was then added, with mixing to form a granulate. The granulate was dried in an Aeromatic Strea-5 fluid bed drier, screened, and mixed with magnesium stearate (100 g). The mixture was filled into hard gelatin capsules to a fill weight of, for example, 500 mg per capsule to achieve a dose of 400 mg of ranolazine dihydrochloride (equivalent to 342 mg of ranolazine free base) per capsule,  
10 but may be filled to fill weight of 30 to 400 mg of ranolazine dihydrochloride.

**EXAMPLE 2**

This Example describes a method of preparing sustained release (SR) ranolazine formulations.

A sustained release (SR) formulation, designated as SR Formulation A, and including pH-dependent and pH-independent binders was prepared by combining Ranolazine (2500 g),  
5 methacrylic acid copolymer, Type C (Eudragit® L 100-55 - Röhm Pharma) (1000 g), microcrystalline cellulose (Avicel®) (100 g) (710 g), and polyvinyl pyrrolidinone powders were intimately mixed together in a Fielder PMA 65 mixer-granulator. The mixture was granulated with a solution of sodium hydroxide (40 g) in water, and a 30% aqueous dispersion of methyl methacrylate/ethyl acrylate copolymer (Eudragit® NE 30 D - Röhm Pharma) (1667  
10 g) was added to the wet mass. The resulting granulate was dried in an Aeromatic Strea-5 fluid bed drier, screened, and then mixed with croscarmellose sodium (100 g) and magnesium stearate (50 g). The mixture was compressed into 684 mg tablets with a Manesty B tablet press to achieve dose of 342 mg of ranolazine free base per tablet. This formulation is referred to as SR Formulation A.

15 SR Formulation B was prepared in the same manner as SR Formulation A except that the Eudragit® L 100-55 was reduced to 500 g, and the Eudragit® NE 30 D was replaced by a 40% aqueous dispersion of a methyl methacrylate/ethyl acrylate copolymer (Eudragit® NE 40 D - Röhm Pharma) (2500 g). The resulting (SR) formulation included 342 mg ranolazine free base per tablet.

20 In SR Formulation C, ranolazine free base (342 mgs) was blended with microcrystalline cellulose and polyvinyl pyrrolininone K25, granulated with water, dried, and blended with croscarmellose sodium and magnesium stearate. The blend was compressed into tablets and coated with an enteric coating.

SR Formulation D, including only a pH dependent binder was prepared by combining  
25 Ranolazine (7500 g), Eudragit® L 100-55 (1000 g), hydroxypropyl methylcellulose

(Methocel® E5 - source) (200 g), and microcrystalline cellulose (Avicel®) (1060 g) by intimate mixing. The mixed powders were granulated with a solution of sodium hydroxide (40 g) in water (1900 to 2500 grams). The granulate was dried and screened, mixed with magnesium stearate (200 g), and compressed for example into tablets weighing 667 mg to achieve a dose of 500 mg of ranolazine free base per tablet. The tablets were spray coated in a 24 inch Accelacota® cylindrical pan coater with OPADRY film coating solution to a 2-4% weight gain. OPADRY film coating solutions are available in a variety of colors from Colorcon, West Point, PA.

The stepwise procedure for preparing SR Formulation D is as follows:

- 10 a) Blend together ranolazine, microcrystalline cellulose, methacrylate copolymer (Type C) and hydroxypropyl methyl cellulose using an appropriate blender.
- b) Dissolve sodium hydroxide in purified water.
- c) Using appropriate granulation equipment, slowly add the sodium hydroxide solution to the blend with constant mixing. Add a further aliquot of water, if necessary.
- 15 d) Continue mixing to achieve additional massing. Add a further aliquot of water, if necessary.
- e) Dry granulated in a fluid bed dryer.
- 20 f) Screen dried granules through an appropriate mill.
- g) Add magnesium stearate to the screened granules and blend together.
- h) Pass the granulated material through a chilsonator, if needed.
- i) Compress the granules into tablets using appropriately sized tooling.
- j) Disperse OPADRY powder in water and film-coat using appropriately sized coating equipment to a typical level of 2-4% by weight.
- 25

**WO 00/13687**

**PCT/US99/20968**

- k) Polish with carnauba wax using a typical level of 0.002-0.003% by weight.**

**EXAMPLE 3**

This Example summarizes a study published in *Circulation* 90:726-734 (1994) that demonstrated that ranolazine was ineffective as an antianginal and anti-ischemic agent when administered as an IR formulation of Example 1.

5        Patients with stable angina pectoris took part in the study. Any previous antianginal drugs used by the patients were discontinued under medical supervision. Three hundred nineteen patients received single-blind placebo for up to 18 days, and 318 stopped exercise because of angina of moderate severity, had evidence of myocardial ischemia ( $\geq$  1-mm ST segment depression), and were randomized to one of four study groups and administered;  
10        ranolazine •2HC1 30 mg tid (n=81); ranolazine •2HC1 60 mg tid (n=81); ranolazine •2HC1 120 mg tid (n=78);, and placebo tid (n=79). After administration of 30-, 60- and 120-mg doses tid, the mean peak plasma concentrations of ranolazine free base at 1 hour after dose were 94, 219 and 510 ng/mL, respectively, and the mean trough plasma concentrations at 8 hours after dose were 18, 37 and 90 ng/mL, respectively.

15        After the 4-week double-blind phase, symptom-limited exercise tests were repeated at 1 hour (peak test) and 8 hours (trough test) after the study medication was administered. Total exercise duration at baseline ( $\pm$ SEM) was  $5.9 \pm 0.2$  minutes for the placebo group and  $6.4 \pm 0.3$ ,  $5.9 \pm 0.3$ , and  $6.6 \pm 0.2$  minutes for the ranolazine 30-, 60-, and 120-mg groups, respectively (P=NS). After 4 weeks of double-blind therapy, compared with baseline values, at 1 hour  
20        after the study medication was administered (peak effect), total exercise duration ( $\pm$ SEM) increased by  $0.45 \pm 0.2$  minutes in the placebo group and by  $0.3 \pm 0.2$ ,  $0.6 \pm 0.2$ , and  $0.5 \pm 0.2$  minutes in the ranolazine 30-, 60- and 120-mg groups, respectively (placebo versus ranolazine, P=NS). Times to 1-mm ST- segment depression at baseline were similar in the  
25        four groups and, after 4 weeks of therapy in each group, increased significantly by similar magnitudes at 1 hour after the administration of the medications. Similar changes were seen

for the time to onset of angina. Eight hours after administration (trough effect), no differences in total exercise time or any other exercise variables were observed between the placebo and the ranolazine groups. Compared with the baseline values, the number of anginal attacks per week and the number and duration of ischemic episodes per 48 hours during Holter  
5 monitoring decreased significantly by similar magnitudes in the placebo and ranolazine groups.

These results indicated that therapy with ranolazine •2HC1 30, 60 and 120 mg tid was not superior to placebo. The study also did not show beneficial effects of similar doses of ranolazine on either myocardial ischemia or exercise performance or on anginal attacks  
10 during daily life in patients with angina pectoris.

**EXAMPLE 4**

In this Example, the safety and anti-ischemic effects of high plasma ranolazine levels in a large group of angina patients was evaluated and the duration of any effects during steady-state dosing with bid and tid regimens was assessed. In this Example, patients with  
5 chronic stable angina pectoris who were responsive to conventional antianginal drugs were treated with 3 ranolazine •2HC1 dosing regimens: 267 mg tid, 400 mg bid and 400 mg tid IR Formulations of Example 1. Exercise testing parameters and ranolazine free base concentrations were determined at peak and trough plasma levels.

**Methods**

10 The study involved double-blind, placebo-controlled randomized treatment phase with 4 treatments (placebo, ranolazine •2HC1 400 mg bid, ranolazine •2HC1 267 mg tid, and ranolazine •2HC1 400 mg tid), 4 treatment sequences and 5 double-blind treatment periods in an extended period Latin square design on pre-qualified patients who were responsive to known antianginal therapy and had stable exercise times.

15 Human patients with chronic stable angina pectoris, of at least 3 months' duration, that had responded to conventional antianginal therapy were considered candidates. In addition, patients had to have electrocardiographic (ECG) evidence of exercise-induced ischemia based upon horizontal or down-sloping ST-segment depression of  $\geq 1$  mm that persisted in 3 consecutive beats during an exercise stress test and an ECG pattern that would not interfere  
20 with interpretation of ST-segment changes. The latter criterion specifically excluded patients with left ventricular hypertrophy, pre-excitation, conduction abnormalities, or pacemaker rhythm. Other exclusion criteria included unstable angina or myocardial infarction within the preceding 3 months, heart failure defined as New York Heart Association Class III or IV, significant valvular or congenital heart disease that was uncorrected, need for digoxin or long-



acting nitrate therapy, labile diabetes mellitus, or other serious conditions that would confuse follow-up evaluation.

These immediate release ranolazine •2HCl dosing regimens (267 mg tid, 400 mg bid, 400 mg tid) and a placebo were administered during the treatment phase. Patients took one capsule containing either 267 mg or 400 mg of ranolazine dihydrochloride, or placebo at 8:00 a.m., 4:00 p.m., 8:00 p.m. and 12:00 a.m. All capsules were identical in appearance. Patients were randomized to 1 of 4 treatment sequences, with 25% of the patients assigned to each sequence. Each treatment was administered for 1 week, with one treatment repeated during a fifth 1-week period.

To qualify patients receiving their usual antianginal medications underwent a screening exercise treadmill test (ETT-1) using a Sheffield modified Bruce protocol. If the time to onset angina was  $\geq 3$  but  $\leq 13$  minutes, an antianginal drug was withdrawn and treatment with single-blind placebo was initiated. After 1 to 2 weeks, patients returned for another ETT (ETT-2). If the time to onset angina decreased by 1 minute compared with ETT-1, the patient was considered to have completed the first qualifying ETT. If the decrease in time to onset angina was not  $\geq 1$  minute, a second antianginal drug could be withdrawn and the above sequence repeated. If necessary, a third antianginal drug could be withdrawn according to this procedure in order for the patient to qualify. Long-acting nitrates were always withdrawn first; beta-blockers; and calcium antagonists could be withdrawn in either order from patients not receiving long-acting nitrates. After the patient achieved the first qualifying ETT (ETT-2), a second qualifying ETT (ETT-3) was performed in which the time to onset angina had to be within  $\pm 15\%$  of that observed during ETT-2. In addition, each of the qualifying ETTs had to have ECG signs of ischemia ( $\geq 1$  mm horizontal or down-sloping ST-segment depression in 3 consecutive beats). Patients meeting these criteria were used in the study.

After each 1 week period, patients returned to the exercise laboratory in the morning, at least 1 hour after a light breakfast, for an ETT. This was designated the trough ETT; the trough ETTs were performed at the same time of day for each patient. After completing the trough ETT, the patient received the next scheduled blinded medication dose from the blister  
5 pack used that week. Another ETT was performed 1 hour after the administered dose. This was designated the peak ETT. Blood samples were obtained at trough (approximately 8 hours after dosing) and at peak (1 hour after dosing). Other standard laboratory tests were monitored regularly throughout the study.

Blood pressure (by cuff) and heart rate were monitored before all ETTs, during the  
10 ETT, during the last minute of each stage of the test, at onset angina, at the point of maximum exercise, and during recovery (every minute for 4 minutes, then every 5 minutes until values returned to baseline). Heart rates also was monitored continuously and standard 12-lead ECG recordings were done immediately before exercise with the patient standing on the treadmill, at the end of each stage of exercise, at the maximally tolerated exercise load, and at the  
15 termination of exercise.

Mean treadmill exercise times for the 3 exercise variables of interest during placebo and the different ranolazine dosing regimens (ranolazine-placebo) for all patients at peak and trough are summarized in Table 2 below.

Table 2

## Exercise Test Data for All Patients at Peak and Trough

Mean Exercise Variable (all patients, minutes)		Treatment			
		Double-blind Placebo	Ranolazine 400 mg bid	Ranolazine 267 mg tid	Ranolazine 400 mg tid
Time to onset angina	Peak	9.01	9.33	9.40	9.33
	Trough	8.58	8.77	8.78	8.65
Exercise Duration	Peak	10.67	10.83	10.87	10.84
	Trough	10.50	10.55	10.56	10.60
Time to 1 mm ST depression	Peak	9.58	9.86	9.99	9.94
	Trough	9.04	9.22	9.22	9.31

5           At peak ranolazine plasma concentrations, all ETT ischemia parameters were prolonged over placebo and most notably, the time to onset of 1-mm ST-segment depression. In the all-patients analysis, the increase in time to onset angina over placebo ranged from 0.32 to 0.39 minutes ( $p \leq 0.01$ ) and time to onset of 1-mm ST-segment depression ranged from 0.28 to 0.41 minutes ( $p \leq 0.02$ ) for each of the 3 ranolazine dosing regimens and all regimens combined. Also, the total duration of exercise was significantly increased for all regimens combined and trends of similar direction and magnitude were noted for each dosing regimen. In the per-protocol analysis each of the 3 ETT parameters were prolonged ( $p \leq 0.01$ ) for all ranolazine dosing regimens combined. All individual ranolazine dosing regimens significantly prolonged time to 1-mm ST-segment depression and nonsignificant trends of

10           similar direction and proportions were found for time to onset angina and duration of exercise. In general, results of the per-protocol analysis, except that the magnitude of the effect appeared to somewhat greater in those with monotherapy.

15           At trough plasma concentrations, ranolazine had a lesser effect on ETT parameters. The results for the all-patients and per-protocol analyses were relatively consistent,

demonstrating trends to increased exercise times. But only the time to 1 mm ST-segment depression for all ranolazine regimens combined in the all-patients analysis achieved statistical significance.

In view of the more pronounced increases in exercise parameters observed with ranolazine monotherapy, the responses to ranolazine among patients receiving different concomitant antianginal medications were analyzed. These post-hoc analyses were performed on peak ranolazine data, when the effects to improve exercise times were most evident. Because long-acting nitrates were withdrawn first during the single-billed qualifying phase, no patient entered double-blind treatment receiving long-acting nitrates. Of patients with peak efficacy data, 34% (107/312) of the patients received beta-blockers during double-blind treatment and 24% (75/312) received calcium antagonists.

Exercise test parameters improved at peak ranolazine concentrations (ranolazine-placebo) whether or not patients were receiving beta-blockers. These improvements were slightly larger in magnitude in the 205 patients not receiving beta-blockers compared to the 107 patients who received beta-blockers. But the differences between those receiving beta-blockers and those not receiving them did not achieve statistical significance for any exercise parameter. In patients not receiving beta-blockers, all exercise parameters improved significantly on each of the 3 ranolazine regimens, and also with all ranolazine regimens combined. Similar trends were observed in the smaller number of patients receiving beta-blockers. Analyses of exercise data from patients taking calcium antagonists compared with those not receiving calcium antagonists produced similar findings.

Table 3 below summarizes the mean peak and trough plasma ranolazine concentrations, in terms of ranolazine dihydrochloride base, for all patients by gender and for each dosing regimen.

Table 3

**Mean ( $\pm$ Standard Deviation) Ranolazine Plasma Concentrations by Dosing Regimen**

	Ranolazine 400 mg bid	Ranolazine 267 mg tid	Ranolazine 400 mg tid
<b>Peak (ng/mL)</b>			
All Patients	1882 (1094)	1346 (823)	2128 (1198)
Males	1762 (999)	1261 (774)	1917 (1046)
Females	2171 (1253)	1594 (904)	2654 (1385)
<b>Trough (ng/mL)</b>			
All Patients	235 (288)	316 (336)	514 (500)
Males	235 (264)	316 (336)	518 (494)
Females	235 (342)	316 (340)	505 (517)

The plasma mean peak concentrations ranged from 1346 to 2128 ng per mL ranolazine free base. The 400 mg tid dosing regimen was associated with the highest ranolazine plasma concentrations. Mean trough ranolazine plasma concentrations ranged from 235 to 514 ng per mL. Mean peak ranolazine plasma concentrations were somewhat higher in females than in males, but there were no sex differences in plasma concentrations evident in trough.

At peak ranolazine plasma concentrations, there were no statistically significant differences among any of the ranolazine dosing regimens and placebo for double product. Likewise, at trough ranolazine plasma concentrations, there were no statistically significant differences among the 3 ranolazine dosing regimens and placebo in the per-protocol analysis for standing or maximum exercise double product.

The results of this study suggest that ranolazine is an effective antianginal and anti-ischemic compound in patients with chronic stable angina pectoris. At peak plasma concentrations, the three ranolazine dosing regimens used prolonged time to onset of angina and duration of exercise as well as time to 1-mm ST-segment depression on average about 0.33 minutes over that observed with placebo. Improvement in exercise parameters was observed in the present study not only in patients receiving concomitant antianginal therapy (eg, beta-blockers and calcium antagonists), but also in the subgroup who received only ranolazine monotherapy. In the latter patients, the treatment effect appeared to be somewhat

greater in magnitude. This suggests that ranolazine also may be useful in monotherapy in patients with chronic stable angina pectoris.

The hemodynamic findings indicate that the improvement in exercise parameters in peak ranolazine plasma concentrations was not associated with changes in blood pressure or heart rate. The nonhemodynamic mechanism of action of ranolazine, therefore, differs from that of other antianginal drugs in current clinical use.

Most important, we documented that the antianginal and anti-ischemic effects of the immediate-release ranolazine preparation studied did not persist throughout the dosing interval. Although time to onset of ischemic-type ST-segment depression was significantly prolonged and trends of similar direction were noted for other ETT parameters, the effect was minimal at trough ranolazine plasma concentrations. Mean peak ranolazine free base plasma concentrations ranged from 1346 to 2128 ng per mL, while mean trough plasma concentrations ranged from 235 to 514 ng per mL. It seems evident that the higher mean ranolazine plasma concentrations observed at peak are associated with clinically meaningful antianginal and anti-ischemic effects, whereas concentrations obtained at trough were not.

Based on the results of the present experiment, the threshold plasma ranolazine free base concentration for anti-ischemic activity detected during ETT is likely to lie above about 550 ng per mL. Further, it is likely that ranolazine plasma concentrations must be maintained at or above the threshold value throughout the dosing interval to ensure antianginal and anti-ischemic activity during exercise throughout this interval.

Ranolazine was well tolerated over the plasma concentrations achieved in the present study. The rate of occurrence of adverse events did not differ among the ranolazine dosing regimens and placebo, and there were no drug-related changes in ECG intervals or complex morphology. In addition, there were no clinically significant changes in blood glucose

concentrations, lipid values or liver function tests, suggesting that the metabolic effect of ranolazine does not extend to systemic glucose regulation or lipid metabolism.

Ranolazine improves exercise parameters with no detectable effect on heart rate and blood pressure in patients with chronic stable angina pectoris. It is likely that a threshold  
5 ranolazine plasma concentration above about 550 ng per mL must be obtained to detect these antianginal and anti-ischemic effects. Ranolazine is well tolerated over a wide range of plasma concentrations. Further study using larger doses of a sustained release preparation are warranted to fully evaluate this novel metabolic concept for management of ischemia.

**EXAMPLE 5****I. *In vitro* Comparison of IR Formulation and SR Formulations**

The IR Formulation prepared according to Example 1 and the SR Formulations prepared according to Examples 2A-2C were tested in a USP Apparatus 2 dissolution tester, using 900 mL of 0.1M hydrochloric acid as the dissolution fluid to simulate dissolution in the stomach.

**Table 4**

Time		Percentage of Formulation Dissolved			
(hours)		Formulation			
		IR	A	B	C
10	0.25	88.1			
	0.5	100.5	13.9	17.6	17.5
	1	101.7	19.9	26.0	25.7
	2		27.8	47.5	35.9
15	4		39.0	69.2	48.4
	8		52.4	90.1	64.7
	12		61.6	99.6	74.2
	24		80.8	105.6	95.4

The tabular results show that while the IR Formulation is completely dissolved in no more than 0.5 hours (as expected for an immediate release formulation), SR Formulations A, B, and C displayed a prolonged dissolution of a low pH, as is desirable for a sustained release formulation.



## II. In vivo Comparison of IR Formulation and SR Formulations A, B, and C

Single doses of the IR Formulation prepared according to Example 1 and SR Formulations A and B prepared according to Example 2 were administered to eleven healthy volunteers and their plasma concentrations of ranolazine free base were measured at 0, 20, 40, 60, 90, and 120 minutes, hourly to six hours, twice-hourly to eighteen hours, and at twenty-four hours after administration (SR Formulations only). The results are set forth in Table 5 below.

Table 5

	Formulation			
	IR	A	B	C
$C_{\max}$ (ISD) (ng/mL)	1940 (807)	753 (264)	657 (316)	925 (747)
$C_{\text{trough}}$ (ISD) (ng/mL)	165 (111)	158 (114)	182 (110)	290 (163)
$T_{\max}$ (ISD) (hours)	1.27 (0.5)	4.09 (1.14)	4.05 (1.31)	6.55 (2.93)
$AUC_{0-24}$ (ISD) (ng.hr/mL)	6530	5640	5280	5820

From Table 5 it is apparent that SR Formulations A, B and C of this invention exhibit dissolution properties which make them suitable for twice daily administration of ranolazine.

**EXAMPLE 6**

This Example details a single-ascending dose, crossover-design study that assessed the safety and pharmacokinetic profile of single oral dose of ranolazine base SR Formulation of Example 2D. Human subjects were divided into three groups. Group 1 received 500, 750 and 1000 mg ranolazine SR. Group 2 received 1250 and 1750 mg ranolazine SR. Group 3 received 1500 and 2000 mg ranolazine SR. Each group also had a randomized placebo phase. Mean pharmacokinetic parameters following single oral doses of the ranolazine SR does are detailed in Table 6 below:

**Table 6**

Mean $\pm$ SD Pharmacokinetic Parameters (n = 8 except* n = 7)					
Dose SR (mg)	Group	C <sub>max</sub> (ng/mL)	C <sub>trough</sub> (ng/mL)	T <sub>max</sub> (hr)	AUC <sub>0-30hr</sub> (ng.hr/mL)
500	1	883 $\pm$ 353	382 $\pm$ 251	4.63 $\pm$ 1.19	9380 $\pm$ 4250
750	1	1300 $\pm$ 1060	455 $\pm$ 353	4.25 $\pm$ 0.886	12500 $\pm$ 9000
1000	1*	1610 $\pm$ 959	695 $\pm$ 438	5.71 $\pm$ 2.14	18100 $\pm$ 9630
1250	2	2210 $\pm$ 716	943 $\pm$ 541	6.21 $\pm$ 3.52	25000 $\pm$ 8090
1500	3	1980 $\pm$ 1050	1070 $\pm$ 754	4.75 $\pm$ 0.886	25400 $\pm$ 16000
1750	2	3670 $\pm$ 1570	2400 $\pm$ 1260	5.25 $\pm$ 2.31	49200 $\pm$ 18200
2000	3	2440 $\pm$ 1120	1640 $\pm$ 937	5.21 $\pm$ 2.30	35400 $\pm$ 19100

10

The pharmacokinetic results reported in Table 6 indicate that ranolazine was slowly released from the SR formulation, and consequently the absorption of ranolazine was dissolution-rate limited. This resulted in prolonged plasma drug concentration-time profiles observed at all dose levels, with peak plasma levels at 4 to 6 hours post dose. Over the dose range 500 to 2000 mg, the mean C<sub>max</sub> and AUC<sub>0-30hr</sub> increased in an approximately dose-proportional manner, although there appeared to be some deviation from proportionality within Group 2.

**EXAMPLE 7**

This Example details a double-blind, placebo-controlled, multiple ascending-dose, crossover-designed volunteer study, to evaluate bid dosing. Six subjects received 4 days dosing with ranolazine SR formulation prepared according to Example 2D at 500, 750, and 5 1000 mg bid, followed by a morning dose on Day 5. Pharmacokinetic results are reported in Table 7, below.

**Table 7**

<b>Day 5 Ranolazine Pharmacokinetic Parameters (mean <math>\pm</math> SD)</b>			
<b>Parameter</b>	<b>Ranolazine SR 500 mg bid (n=7)</b>	<b>Ranolazine SR 750 mg bid (n=7)</b>	<b>Ranolazine SR 1000 mg bid (n=7)</b>
$C_{max}$ (ng/mL)	1760 $\pm$ 715	2710 $\pm$ 657	3660 $\pm$ 1090
$T_{max}$ (hr)	2.00 $\pm$ 1.15	4.33 $\pm$ 1.62	4.17 $\pm$ 2.48
$C_{min}$ (ng/mL)	585 $\pm$ 340	1260 $\pm$ 501	1960 $\pm$ 812

According to Table 7, ranolazine was slowly released from the SR formulation, and 10 consequently the pharmacokinetics were dissolution-rate limited. This resulted in extended plasma drug concentration-time profiles at all dose levels, with peak plasma levels observed at 2 to 4 hours post dose.

These results indicate that useful ranolazine plasma levels can be achieved in humans with dosing of this SR formulation on a bid schedule.

**EXAMPLE 8**

This Example evaluated the safety and tolerability of administering racemic ranolazine free base formulations as in Example 2D. The individual and mean concentrations of racemic ranolazine and its enantiomers, (R)-(+)-N-(2,6-dimethylphenyl)-4-[2-hydroxy-3-(2-methoxyphenoxy)-propyl]-1-piperazineacetamide, (S) - (-) - N - (2,6-dimethylphenyl) -4-[2-hydroxy-3-(2-methoxyphenoxy)-propyl]-1-piperazineacetamide in human plasma were also determined.

The study was performed with ascending doses of sustained release ranolazine dosage forms. Before and at intervals during and after the dosing period, blood samples were drawn for ranolazine assay and blood pressure, heart rate, ECG and symptoms were monitored throughout. Data summaries were reviewed after each phase before proceeding to the next phase of the study.

Eight subjects, all healthy male volunteers aged between 18 and 40 entered and all completed the study and were available for pharmacokinetic and safety analysis. The subjects were each given doses of the various types of ranolazine free base in the form of sustained release tablets including 500 mg and 750 mg tablets or with matching placebo as necessary (2 x 750 mg size plus 1 x 500 mg size) to make unit oral doses of 1500 and 2000 mg.

In each phase: one dose bid for four days with a single dose on Day 5. On day 5 each volunteer underwent a full pharmacokinetic profile including supine and erect blood pressure (BP) and heart rate, ECG data, adverse events, clinical chemistry and hematology results, urinalysis results.

Steady state was tested for in each dose level, using  $C_{48h}$ ,  $C_{72h}$  and  $C_{96h}$  and log transformed data, by analysis of covariance and by testing whether the coefficient for time was significantly different (defined as  $p < 0.05$ ) from 0. These tests were made using two-sided t-tests with estimates of variability from the ANOVA models. Steady state was also

assessed by comparing means for  $C_{48h}$ ,  $C_{72h}$  and  $C_{96h}$  using a mixed effects ANOVA model and untransformed and log, transformed data. For hemodynamic parameters, Day 1 pre-dose treatment means and Day 5 data were compared across treatments via two-sided t-tests using estimates of variability from mixed effects ANOVA models. Ninety and 95% confidence intervals were calculated for the treatment comparisons. No adjustments were made for multiple comparisons.

The mean and standard deviation day 5 pharmacokinetic parameters of ranolazine free base are detailed in Table 8 below and the mean plasma profiles are shown in the figure. Steady-state plasma levels of ranolazine free base appeared to be attained by day 4. Within the dose interval there was a slow rise to maximum levels with  $t_{max}$  values ranging from 1 to 6 h post-dose. Thereafter levels declined slowly, producing a small degree of fluctuation in plasma levels over the dosing interval. There appeared to be no differences in the pharmacokinetic parameters of the (+) R and (-) S enantiomers of ranolazine following multiple dosing with this SR formulation.

15

**Table 8**  
**Day 5 Racemic (RS), (+)R and (-)S Ranolazine SR Pharmacokinetic Parameters**

Parameter	1500 mg SR bid			2000 mg SR bid
	RS ranolazine	(+)R ranolazine	(-)S ranolazine	RS ranolazine
$C_{max}$ (ng/ml)	5284 ± 2434	2909 ± 1308	2944 ± 1426	7281 ± 2700
$C_{min}$ (ng/ml)	2932 ± 1918	1436 ± 1046	1514 ± 1201	4149 ± 2228
Median $t_{max}$ (h)	4.00	4.00	4.00	4.00
$C_{96h}$ (ng/ml)	3656 ± 1918	2009 ± 1137	2399 ± 1205	5042 ± 1797
$C_{108h}$ (ng/ml)	2942 ± 1937	1447 ± 1071	1541 ± 1260	4398 ± 2396
$AUC_{96-108h}$ (ng.h/ml)	49516 ± 23945	25731 ± 13385	26407 ± 14849	68459 ± 25842
$C_{ave}$ (ng/ml)	4126 ± 1995	2144 ± 1115	2201 ± 1237	5705 ± 2153
Degree of fluctuation	0.664 ± 0.336	0.833 ± 0.402	0.824 ± 0.443	0.591 ± 0.240

Some subjects became too symptomatic on standing to complete BP measurements on ranolazine 1500 mg (n=3 of 8) and 2000 mg (n=2 of 8), usually 2-6 hours post-dose. Statistically significant reductions in orthostatic systolic BP were noted on day 5 on 1500 mg

(-9.8 mm Hg; 4 hours post-dose) and 2000 mg (-8.4 mmHg; 6 hours post-dose). Although the pattern of adverse events was similar on ranolazine and placebo, headache, dizziness and blocked nose seemed more common on ranolazine.

What we claim is:

1. A method for treating a human patient suffering from a cardiovascular disease selected from arrhythmias, variant and exercise-induced angina, and myocardial infarction by administering a sustained release pharmaceutical dosage form including at least 50% by weight ranolazine in no more than two tablets per dose to the human patient to maintain ranolazine plasma levels in the human patient of from about 550 to about 7500 ng base/mL for at least 24 hours wherein the dose is administered at a frequency selected from once, twice and three times over 24 hours.
2. The method of claim 1 wherein the sustained release dosage form includes at least one pH dependent binder wherein the pH dependent binder inhibits the release of ranolazine from the sustained release dosage form when the sustained release dosage form is subjected to an aqueous environment having a pH of the stomach and wherein the pH dependent binder promotes the release of a therapeutic amount of ranolazine in an aqueous solution having a pH above about 4.5.
3. The method of claim 2 wherein the pH dependent binder is partially neutralized.
4. The method of claim 1 wherein the pharmaceutical dosage form is administered to the human patient at a frequency selected from once and twice over 24 hours.
5. The method of claim 1 wherein the pharmaceutical dosage form is administered to the human patient in two doses over 24 hours wherein each dose consists of two tablets.
6. The method of claim 1 wherein the pharmaceutical dosage form includes between about 50% to about 95% by weight ranolazine.
7. The method of claim 1 wherein the pharmaceutical dosage form includes from about 70% to about 80% by weight ranolazine.
8. The method of claim 2 wherein the pH dependent binder is selected from methacrylic acid copolymers, hydroxypropyl cellulose phthalate, hydroxypropyl methylcellulose

phthalate, cellulose acetate phthalate, polyvinyl acetate, phthalate, polyvinylpyrrolidine phthalate, and mixtures thereof.

9. The method of claim 2 wherein the pH dependent binder is a methacrylic acid copolymer.
10. The method of claim 9 wherein the methacrylic acid copolymer is methacrylic acid  
5 copolymer Type C USP.
11. The method of claim 2 wherein the pH dependent binder is from about 5 to about 12 wt% methacrylic acid copolymer Type C USP.
12. The method of claim 1 wherein the pH dependent binder is about 10wt% methacrylic acid copolymer Type C USP.
- 10 13. The method of claim 1 wherein the pharmaceutical dosage form includes a pH-independent binder.
14. The method of claim 13 wherein the pH-independent binder is selected from hydroxypropyl methylcellulose, hydroxypropyl cellulose, poly(meth)acrylate esters, polyvinylpyrrolidone, and mixtures thereof.
- 15 15. The method of claim 13 wherein the pH-independent binder is hydroxypropyl methylcellulose.
16. The method of claim 15 wherein the pharmaceutical dosage form includes from about 1 to about 3 wt% hydroxypropyl methylcellulose.
17. The method of claim 15 wherein the pharmaceutical dosage form includes about 2 wt%  
20 hydroxypropyl methylcellulose.
18. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 1000-5000 ng base/mL.
19. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 1000-3800 ng base/mL.



20. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 550-5000 ng base/mL.
21. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 550-3800 ng base/mL.
- 5 22. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 1000-2800 ng base/mL.
23. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 1700-3900 ng base/mL.
24. The method of claim 1 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 wherein the human patient plasma ranolazine level ranges from 550-2000 ng base/mL.
- 10 25. The method of claim 23 wherein the dosage form includes from 650-850 mg ranolazine.
26. The method of claim 24 wherein the dosage form includes from 900-1100 mg ranolazine.
27. The method of claim 25 wherein the dosage form includes from 400-600 mg ranolazine.
28. The method of claim 22 or 23 or 24 or 25 or 26 or 27 wherein the peak to trough human patient plasma ranolazine levels is less than 4:1 over a 24 hour period.
- 15 29. The method of claim 23 or 24 or 25 or 26 or 27 wherein the peak to trough human patient plasma ranolazine levels is less than 3:1 over a 24 hour period.
30. The method of claim 24 or 28 wherein the peak to trough human patient plasma ranolazine levels is less than 2:1 over a 24 hour period.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 31/495, 9/20</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 00/13687</b> <b>(43) International Publication Date:</b> 16 March 2000 (16.03.00)
<b>(21) International Application Number:</b> PCT/US99/20968 <b>(22) International Filing Date:</b> 9 September 1999 (09.09.99)  <b>(30) Priority Data:</b> 60/099,804 10 September 1998 (10.09.98) US 09/321,522 27 May 1999 (27.05.99) US  <b>(71) Applicant (for all designated States except US):</b> CV THERAPEUTICS, INC. [US/US]; 3172 Porter Drive, Palo Alto, CA 94304 (US).  <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> WOLFF, Andrew, A. [US/US]; 1140 Lawton Street, San Francisco, CA 94122 (US).  <b>(74) Agent:</b> HUGHES, A., Blair, McDonnell, Boehnen, Hulbert & Berghoff, Suite 3200, 300 S. Wacker Drive, Chicago, IL 60606 (US).	<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 29 June 2000 (29.06.00)	
<b>(54) Title:</b> SUSTAINED RELEASE RANOLAZINE FORMULATIONS  <b>(57) Abstract</b>  A sustained release ranolazine formulation contains an intimate mixture of ranolazine and a partially neutralized pH-dependent binder to form a film that is mostly insoluble in aqueous media below pH 4.5 and soluble in aqueous media above pH 4.5. The formulation is suitable for twice daily administration of ranolazine and is useful for controlling the rate of dissolution of ranolazine, and to maintain human plasma ranolazine levels at between 550 and 7500 ng base/mL.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

Intern. Application No.  
PCT/US 99/20968

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K31/495 A61K9/20

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 26266 A (SYNTEX) 24 November 1994 (1994-11-24) cited in the application claims 1,2,4-6,10-13,15 page 3, line 16 - line 19 page 4, line 37 -page 5, line 13 page 7, line 9 - line 11	1,2,6,8, 13,20,24
A	EP 0 407 780 A (SYNTEX) 16 January 1991 (1991-01-16) cited in the application claims 14,15 examples C,D	1,2,6,8, 13,20,24

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

15 March 2000

Date of mailing of the international search report

27/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Scarponi, U

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/20968

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 567 264 A (A.F.KLUGE ET AL.) 28 January 1986 (1986-01-28) cited in the application claims examples 12,14	1,2,6,8, 13,20,24
A	CHEMICAL ABSTRACTS, vol. 121, no. 23, 5 December 1994 (1994-12-05) Columbus, Ohio, US; abstract no. 271766, XP002133147 abstract & M.R.GRALINSKI ET AL.: "CARDIOPROTECTIVE EFFECTS OF RANOLAZINE (RS-43285) IN THE ISOLATED PERFUSED RABBIT HEART" CARDIOVASC. RES., vol. 28, no. 8, 1994, pages 1231-1237,	1,2,6,8, 13,20,24
A	CHEMICAL ABSTRACTS, vol. 129, no. 1, 6 July 1998 (1998-07-06) Columbus, Ohio, US; abstract no. 24, XP002133148 abstract & J.G.MCCORMACK ET AL.: "RANOLAZINE:A NOVEL METABOLIC MODULATOR FOR THE TREATMENT OF ANGINA." GEN. PHARMACOL., vol. 30, no. 5, 1998, pages 639-645,	1,2,6,8, 13,20,24

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 20968

### Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
**Remark: Although claims 1-30 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.**
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No  
PCT/US 99/20968

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9426266 A	24-11-1994	US 5455045 A	03-10-1995
		AT 187067 T	15-12-1999
		AU 674960 B	16-01-1997
		AU 6830194 A	12-12-1994
		BR 9406604 A	02-01-1996
		CA 2162670 A	24-11-1994
		CN 1122571 A, B	15-05-1996
		CZ 9502954 A	14-02-1996
		DE 69421918 D	05-01-2000
		EP 0697866 A	28-02-1996
		FI 955439 A	10-11-1995
		HU 9500110 A	28-06-1995
		HU 72978 A	28-06-1996
		IL 109639 A	08-02-1998
		JP 8510254 T	29-10-1996
		NO 954546 A	10-11-1995
		NZ 266683 A	27-08-1996
		PL 311654 A	04-03-1996
		US 5472707 A	05-12-1995
		US 5554384 A	10-09-1996
EP 407780 A	16-01-1991	AT 142497 T	15-09-1996
		AU 633589 B	04-02-1993
		AU 5761890 A	03-01-1991
		CA 2019580 A	23-12-1990
		DE 69028457 D	17-10-1996
		DK 407780 T	30-09-1996
		EP 0714660 A	05-06-1996
		EP 0719558 A	03-07-1996
		ES 2091211 T	01-11-1996
		GR 3020976 T	31-12-1996
		IE 80710 B	16-12-1998
		JP 3048672 A	01-03-1991
		NZ 234184 A	29-01-1997
		NZ 247044 A	24-04-1997
		US 5906988 A	25-05-1999
		US 5506229 A	09-04-1996
		ZA 9004842 A	26-02-1992
US 4567264 A	28-01-1986	AT 31533 T	15-01-1988
		AU 566489 B	22-10-1987
		AU 2834684 A	22-11-1984
		CA 1256874 A	04-07-1989
		CS 246099 B	16-10-1986
		CS 246080 B	16-10-1986
		DK 248384 A	19-11-1984
		EP 0126449 A	28-11-1984
		ES 532565 D	01-12-1985
		ES 8601947 A	01-03-1986
		FI 841989 A, B,	19-11-1984
		HK 91989 A	24-11-1989
		HU 34177 A, B	28-02-1985
		IE 57487 B	10-02-1993
		IL 71863 A	30-10-1987
		JP 1780695 C	13-08-1993
		JP 4069151 B	05-11-1992
		JP 59219271 A	10-12-1984
		KR 8900020 B	06-03-1989

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/20968

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4567264 A		MX 9203001 A	01-07-1992
		NO 841968 A,B,	19-11-1984
		NZ 208188 A	30-10-1987
		PH 20016 A	01-09-1986
		PL 247722 A	13-08-1985
		PL 252856 A	19-11-1985
		PT 78604 A,B	01-06-1984
		SG 59889 G	29-12-1989
		RU 2071471 C	10-01-1997
		RU 2083570 C	10-07-1997
		ZA 8403746 A	29-01-1986